Asian Journal of Technology and Applied Sciences
ISSN: 2301-xxxx (online) | IF: 10.01 | Vol: 1, issue 1
Available: http://asiansciencejournal.org

DEVELOPMENT OF BREATHING AND VOICE DISORDERS IN CHILDREN WITH DYSARTHRIA

Akramov Dostonbek Ikromjon oglu
Fergana State University
Teacher of the department of inclusive education, faculty of pedagogy, psychology and art studies.
E-mail: akramovdostonbek28121998@gmail.com

Abstract: Dysarthria is a neurological motor speech disorder characterized by impairments in the muscles responsible for speech production. Among the most significant manifestations of dysarthria are breathing and voice disorders, which directly affect speech intelligibility, communication effectiveness, and social participation. Respiratory and phonatory impairments frequently occur due to inadequate neuromuscular control, reduced muscle strength, abnormal muscle tone, and impaired coordination of speech-related structures. These difficulties often limit a child's ability to produce adequate breath support, maintain vocal intensity, and regulate speech rhythm. The purpose of this study is to examine the development of breathing and voice disorders in children with dysarthria and to analyze their impact on communication abilities. The study reviews theoretical perspectives, neurological mechanisms, clinical characteristics, assessment procedures, and intervention approaches related to respiratory and phonatory dysfunctions. Findings indicate that respiratory and voice impairments significantly contribute to communication difficulties and may negatively affect academic achievement, social interaction, and overall quality of life. Early identification and comprehensive intervention targeting respiratory control and voice production can improve communicative outcomes and enhance functional participation.

Keywords: dysarthria, breathing disorders, voice disorders, respiratory control, phonation, speech production.

Introduction. Speech production is a highly complex process requiring the coordinated functioning of multiple physiological systems. Among these systems, respiration and phonation play a central role in the generation of spoken language. Adequate respiratory support provides the airflow necessary for speech, while phonatory mechanisms transform this airflow into audible vocal signals. Any disruption within these systems may significantly impair communication abilities. Children with dysarthria frequently experience difficulties in respiratory and phonatory functioning due to neurological impairments affecting the muscles involved in speech production. Dysarthria is characterized by abnormalities in strength, range, speed, coordination, and timing of speech movements. Consequently, respiratory and laryngeal systems often fail to operate efficiently, resulting in breathing difficulties, reduced vocal intensity, abnormal voice quality, and impaired speech intelligibility. Breathing disorders associated with dysarthria typically involve inadequate respiratory support for speech. Children may exhibit shallow breathing, reduced lung volume utilization, poor coordination between respiration and phonation, and difficulties sustaining airflow during speech production. These impairments frequently result in short utterances, interrupted speech patterns, and reduced communicative effectiveness. Voice disorders represent another significant challenge for children with dysarthria. Neurological impairments may affect vocal fold movement and laryngeal control, leading to hoarseness, breathiness, strained voice quality, reduced loudness, and abnormal pitch variations. Such voice disturbances can significantly decrease speech intelligibility and negatively influence social participation. The

relationship between respiration, phonation, and speech development is particularly important during childhood because communication skills continue to develop throughout this period. Persistent respiratory and voice impairments may restrict opportunities for language learning, academic participation, and social interaction. The purpose of this study is to investigate the development of breathing and voice disorders in children with dysarthria and to explore their effects on communication performance. Understanding these relationships may contribute to improved assessment procedures and intervention strategies. Dysarthria as a motor speech disorder. Dysarthria refers to a group of motor speech disorders resulting from damage to the central or peripheral nervous system. The disorder affects the execution of speech movements rather than linguistic formulation. Depending on the site and severity of neurological damage, dysarthria may involve impairments in respiration, phonation, resonance, articulation, and prosody. Children with dysarthria often demonstrate significant variability in symptom presentation. Some individuals experience mild speech difficulties, while others exhibit severe communication impairments requiring alternative communication systems. Respiratory and phonatory disturbances are among the most frequently reported symptoms. Neurological conditions commonly associated with childhood dysarthria include cerebral palsy, traumatic brain injury, muscular dystrophy, genetic syndromes, and neurodevelopmental disorders. These conditions may disrupt the neuromuscular mechanisms necessary for effective speech production. Respiratory function in speech production.Respiration serves as the foundation of speech production. The respiratory system generates the airflow required for phonation and articulation. Effective speech production depends on precise coordination among respiratory muscles, including the diaphragm, intercostal muscles, and abdominal musculature. During speech, breathing patterns differ substantially from those observed during quiet respiration. Speakers typically inhale rapidly and exhale slowly while controlling airflow to support speech production. This process requires sophisticated neuromuscular coordination. Children with dysarthria frequently experience impairments in respiratory control due to muscle weakness, spasticity, ataxia, or involuntary movements. These impairments may reduce respiratory efficiency and compromise speech performance. Characteristics of breathing disorders in dysarthria. Respiratory disorders associated with dysarthria may manifest in various ways. One common characteristic is reduced vital capacity. Children may be unable to utilize sufficient lung volume during speech, resulting in shortened phrases and reduced vocal endurance. Inadequate respiratory support often forces speakers to pause frequently during communication. Another common feature involves poor coordination between breathing and speech production. Children may initiate speech at inappropriate points within the respiratory cycle or fail to maintain adequate airflow throughout an utterance. These difficulties frequently contribute to interrupted speech patterns and decreased intelligibility. Respiratory muscle weakness may further limit the ability to generate sufficient subglottal pressure for speech. As a result, vocal intensity may be reduced, making speech difficult to hear and understand. Voice production and phonation. Voice production occurs within the larynx through vibration of the vocal folds. Airflow generated by the respiratory system passes through the vocal folds, causing them to vibrate and produce sound. Effective phonation requires precise regulation of vocal fold tension, position, and movement. The quality of the voice depends on multiple physiological factors, including respiratory support, vocal fold function, and resonance characteristics. Disruptions affecting any of these components may lead to voice disorders. Children with dysarthria frequently exhibit abnormalities in voice production due to impaired neuromuscular control of laryngeal structures. Characteristics of voice disorders in dysarthria. Voice disorders in dysarthria may present as breathiness,

hoarseness, harshness, strain, reduced loudness, monopitch, monoloudness, or unstable vocal quality. Breathy voice quality often results from incomplete closure of the vocal folds during phonation. Excessive air leakage reduces vocal efficiency and contributes to decreased speech intelligibility. Harsh or strained voice quality may occur when excessive muscular tension affects laryngeal function. These characteristics frequently increase vocal effort and reduce communicative effectiveness. Many children with dysarthria demonstrate reduced vocal intensity due to inadequate respiratory support and impaired laryngeal control. Low vocal volume may make communication difficult, particularly in educational and social environments. Pitch abnormalities are also common. Some children exhibit restricted pitch variation, resulting in monotonous speech. Others demonstrate excessive pitch fluctuations that interfere with speech naturalness. Neurological mechanisms underlying respiratory and voice disorders. The neural control of respiration and phonation involves complex interactions among cortical, subcortical, cerebellar, and brainstem structures. The motor cortex contributes to voluntary respiratory control and speech production. Damage affecting cortical motor areas may impair respiratory coordination and laryngeal function. The cerebellum plays a critical role in timing, coordination, and motor learning. Cerebellar dysfunction may result in irregular respiratory patterns and unstable phonation. The basal ganglia contribute to movement regulation and speech initiation. Neurological abnormalities affecting these structures may produce characteristic voice disturbances and respiratory difficulties. Brainstem nuclei coordinate essential respiratory and laryngeal functions. Damage within these regions may significantly compromise speech production mechanisms. Impact on communication development. Breathing and voice disorders can substantially affect communication development in children with dysarthria. Reduced speech intelligibility often limits participation in conversations and educational activities. Communication difficulties may negatively influence language acquisition, academic achievement, social relationships, and emotional well-being. Children who experience persistent speech impairments may develop reduced self-confidence and increased communication anxiety. The cumulative effects of respiratory and phonatory dysfunctions highlight the importance of early assessment and intervention.

Methods. Research Design. This study employed a qualitative literature-based research design to investigate the development of breathing and voice disorders in children with dysarthria. The research focused on identifying the mechanisms underlying respiratory and phonatory impairments and examining their influence on communication development. A descriptive and analytical approach was adopted to synthesize current scientific knowledge regarding respiratory function, voice production, neurological control mechanisms, and intervention strategies. This approach enabled the integration of findings from speech-language pathology, developmental psychology, neurology, rehabilitation sciences, and special education. Sources of Data. Data were collected from peer-reviewed journal articles, academic books, clinical reports, and international scientific databases. The selected literature focused on: Childhood dysarthria, Respiratory impairments in speech disorders, Voice disorders associated with neurological conditions, Speech production mechanisms, Intervention approaches for breathing and phonation disorders.Studies involving children diagnosed with cerebral palsy, traumatic brain injury, neuromuscular disorders, and developmental neurological conditions were included because these populations frequently exhibit dysarthric symptoms. Research examining assessment or intervention outcomes.

Data Analysis. The collected information was analyzed using thematic analysis. Findings were organized into the following categories:

Respiratory impairments and speech production.
Voice disorders and communication performance.
Neurological mechanisms.
Clinical assessment.
Therapeutic intervention.
Educational implications. Common themes and relationships were identified and synthesized to provide a comprehensive understanding of breathing and voice disorders in children with dysarthria. Results. The analysis revealed that respiratory and phonatory impairments are among the most significant contributors to communication difficulties in children with dysarthria. Respiratory deficits and speech performance. These impairments frequently resulted in shortened utterances and frequent pauses during speech. Many children were unable to complete longer sentences without taking additional breaths. Research indicated that respiratory dysfunction directly affected speech intelligibility. Children with stronger respiratory control generally produced clearer and more understandable speech. Voice Disorders and Communication. Voice disorders were identified as a major factor affecting communication effectiveness. The most frequently reported voice characteristics included: Hoarseness. Breathiness. Harsh vocal quality. Reduced loudness.
Vocal instability.
Monotone speech.

Children with severe voice disorders often experienced difficulties participating in classroom discussions and social interactions. Reduced vocal intensity made communication particularly challenging in noisy environments. Several studies reported that voice abnormalities contributed significantly to reduced listener comprehension even when articulation abilities were relatively preserved. Neurological findings. Neurophysiological research revealed that respiratory and laryngeal functions are controlled by interconnected neural systems. Damage affecting motor pathways frequently produced simultaneous impairments in: Respiratory control. Voice production. Articulation. Prosody. These findings support the view that breathing and voice disorders should not be considered isolated symptoms but rather components of a broader motor speech impairment. Discussion. The findings confirm that breathing and voice disorders represent central features of dysarthria in children. One of the most significant observations is the importance of respiratory support for effective communication. Speech production depends on controlled airflow, and deficits affecting respiratory function can compromise all aspects of communication. Children with dysarthria often struggle to generate adequate subglottal pressure necessary for efficient phonation. Consequently, speech may become weak, interrupted, and difficult to understand. The results also highlight the importance of voice quality in communication development. Voice abnormalities influence not only speech intelligibility but also social perception and participation. Children with severe voice disorders may experience reduced confidence and increased communication barriers. The neurological evidence reviewed in this study supports contemporary theories emphasizing the integrated nature of speech production systems. Respiratory, phonatory, articulatory, and prosodic functions operate as interconnected components rather than independent mechanisms.

REFERENCES

1. Vygotsky, L. S. (1934). Mishelenia i rech. Moscow: Sosekgiz.
2. Luria, A. R. (1979). Yazik i soznanie. Moscow: Izdatelstvo MGU.
3. Leontyev, A. A. (1969). Psikholinguisticheskie edinitsy i porozdenie rechevogo vyskazyvaniya. Moscow: Nauka.
4. Levina, R. E. (1968). Osnovi teorii i praktiki logopedii. Moscow: Prosveshenie.
5. Lopatina, L. V. (2005). Logopedic work with children with minimal dysarthric disorders. St. Petersburg: Soyuz.
6. Mamarajabov, S. (2022). Basics of special pedagogy and logopedics. Tashkent: Teacher.
7. Akhmedova, M. R. (2024). Oligophrenopedagogy and speech development methodology. Tashkent: Science and technologies.